

*Deltocephalus pulicaris* (Fall.) one male, two females.  
*Psammotettix confinis* (Dahlb.) three males (and four *Psammotettix* females).  
*Psammotettix nodosus* (Ribaut) three males.  
*Elymana sulphurella* (Zett.) two females.  
*Conomelus anceps* (Germ.) many.  
*Sardius argus* (Marsh.) one female.

#### LEPIDOPTERA ON LUNDY 1995

The figures given are the earliest and latest sightings, followed by a figure in brackets which is the maximum recorded in any one day.

Orange Swift *Hepialus sylvina* (L.) 30.7 - 4.8.95 (6)  
 Five-spot Burnet *Zygaena trifolii* (Esp.) larvae 25.4 - 3.6.95 (6); moths 28.6 - 19.7.95 (10)  
*Monopis rusticella* (Hbn.) 7.6.95 (several) in dead Soay sheep under Logan Stone (R.S.Key, determined R.Johnson)  
*Ypsolopha dentella* (Fab.) 3.8.95 (1)  
*Plutella xylostella* (L.) 30.7 - 4.8.95 (10)  
*Hofmannophila pseudospretella* (Staint.) 1.8.95 (1)  
*Depressaria pastinacella* (Dup.) 4.8.95 (1)  
*Agonopterix subpropinquella* (Staint.) 1.8 - 4.8.95 (5)  
*Agonopterix alstroemeriana* (Clerck) 29.7.95 (1)  
*Rhyacionia buoliana* (D. & S.) 29.7 - 3.8.95 (6)  
*Epiblema uddmanniana* (L.) 29.7 - 30.7.95 (2)  
*Epiblema roborana* (D. & S.) 3.8.95 (2)  
*Bactra lancealana* (Hbn.) 31.7.95 (5)  
*Olethreutes lacunana* (D. & S.) 1.8.95 (1)  
*Pandemis cerasana* (Hbn.) 1.8 - 2.8.95 (1)  
*Archips podana* (Scop.) 3.8.95 (1)  
*Croesia forsskaleana* (L.) 29.7 - 2.8.95 (9)  
*Agapeta hamana* (L.) 1.8 - 3.8.95 (4)  
*Aethes rubigana* (Treit.) 2.8.95 (1)  
*Alucita hexadactyla* (L.) 29.7 - 2.8.95 (2)  
*Agriphila straminella* (D. & S.) 29.7 - 4.8.95 (20)  
*Agriphila tristella* (D. & S.) 30.7 - 1.8.95 (4)  
*Agriphila geniculata* (Haw.) 4.8.95 (1)  
*Catoptria pinella* (L.) 1.8.95 (1) St John's Valley  
*Catoptria margaritella* (D. & S.) 1.8.95 (1) St John's Valley  
*Dipleurina lacustrata* (Panz.) 29.7 - 2.8.95 (7)  
*Eudonia mercurella* (L.) 29.7 - 4.8.95 (3)  
*Pyrausta cespitalis* (D. & S.) 30.7.95 (1)  
*Udea ferrugalis* (Hbn.) 29.7.95 (3)  
*Nomophila noctuella* (D. & S.) 29.7.95 (2)  
*Dolicharthria punctalis* (D. & S.) 29.7 - 3.8.95 (2)  
*Pleuroptya ruralis* (Scop.) 29.7.95 (10)  
*Endotrichia flammealis* (D. & S.) 29.7.95 (1)  
 Large Skipper *Ochloides venata* (Brem. & Grey) 9.6.95 (1) above Quarry Bay (R.S.Key); new record.  
 Clouded Yellow *Colias croceus* (Geoff.) 4.8.95 (1)  
 Large White *Pieris brassicae* (L.) 4.5 - 21.9.95 (60)  
 Small White *Pieris rapae* (L.) 2.5 - 30.9.95 (500)  
 Green-veined White *Pieris napi* (L.) 6.6 - 30.8.95 (3)  
 Small Copper *Lycaena phlaeas* (Fab.) 8.5 - 13.10.95 (9); 17.8.95, a female blue-spotted aberration  
 Common Blue *Polyommatus icarus* (Rott.) 4.8 - 19.8.95 (1)  
 Red Admiral *Vanessa atalanta* (L.) 25.3 - 13.11.95 (100)  
 Painted Lady *Cynthia cardui* (L.) 6.5 - 27.10.95 (50)  
 Small Tortoiseshell *Aglais urticae* (L.) 4.4 - 13.10.95 (12)  
 Peacock *Inachis io* (L.) 27.7 - 13.8.95 (2)

Speckled Wood *Pararge aegeria* (L.) 15.6 - 18.8.95 (1)  
 Wall *Lasionmata megera* (L.) 24.6 - 8.8.95 (6)  
 Grayling *Hipparchia semele* (L.) 6.5 - 1.9.95 (30)  
 Gatekeeper *Pyronia tithonus* (L.) 25.6 - 3.8.95 (4)  
 Meadow Brown *Maniola jurtina* (L.) 15.6 - 19.8.95 (100)  
 Small Heath *Coenonympha pamphilus* (L.) 31.5 - 17.8.95 (15)  
 Ringlet *Aphantopus hyperantus* (L.) 1.7.95 (1)  
 Fox Moth *Macrothylacia rubi* (L.) larvae 30.5 - 15.8.95 (2)  
 Emperor Moth *Pavonia pavonia* (L.) 6.5.96 (1)  
 Chinese Character *Cilix glaucata* (Scop.) 29.7 - 4.8.95 (3)  
 Mullein Wave *Scopula marginepunctata* (Goeze) 3.8 - 4.8.95 (3)  
 Small Blood-vein *Scopula imitaria* (Hbn.) 29.7 - 1.8.95 (1)  
 Lesser Cream Wave *Scopula immutata* (L.) 1.8 - 3.8.95 (1)  
 Small Fan-footed Wave *Idaea biselata* (Hufn.) 29.7 - 4.8.95 (8)  
 Single-dotted Wave *Idaea dimidiata* (Hufn.) 2.8.95 (1)  
 Riband Wave *Idaea aversata* (L.) 1.8.95 (3)  
 Garden Carpet *Xanthorhoe fluctuata* (L.) 30.7 - 3.8.95 (1)  
 Red Twin-spot Carpet *Xanthorhoe spadicearia* (D. & S.) 30.7 - 1.8.95 (2)  
 Common Carpet *Epirrhoe alternata* (Mll.) 3.8.95 (1)  
 Yellow Shell *Campptogramma bilineata* (L.) 29.7 - 4.8.95 (4)  
 Small Phoenix *Ecliptoptera silaceata* (D. & S.) 29.7.95 (1)  
 July Highflyer *Hydriomena furcata* (Thunb.) 3.8.95 (1)  
 Rivulet *Perizoma affinitata* (Steph.) 29.7 - 2.8.95 (8)  
 Small Rivulet *Perizoma alchemillata* (L.) 29.7 - 3.8.95 (3)  
 Slender Pug *Eupithecia tenuiata* (Hbn.) 29.7.95 (1)  
 Foxglove Pug *Eupithecia pulchellata* Steph. 29.7.95 (5)  
 Bleached Pug *Eupithecia expallidata* Doubl. 29.7 - 2.8.95 (2)  
 Green Pug *Chloroclystis rectangulata* (L.) 1.8.95 (1)  
 Maggie Moth *Abraxas grossulariata* (L.) 29.7 - 14.8.95 (7)  
 Brimstone Moth *Opisthograptis luteolata* (L.) 1.8 - 4.8.95 (7)  
 Early Thorn *Selenia dentaria* (Fab.) 1.8 - 3.8.95 (1)  
 Purple Thorn *Selenia tetralunaria* (Hufn.) 30.7.95 (1)  
 Scalloped Oak *Crocallis elinguaris* (L.) 29.7 - 1.8.95 (3)  
 Peppered *Biston betularia* (L.) 29.7 - 2.8.95 (4)  
 Willow Beauty *Peribatodes rhomboidaria* (D. & S.) 29.7 - 2.8.95 (4)  
 Mottles Beauty *Alcis repandata* (L.) 1.8.95 (1)  
 Engrailed *Ectropis bistortata* (Goeze) 3.8.95 (1)  
 Light Emerald *Campaea margaritata* (L.) 1.8.95 (1)  
 Humming-bird Hawk-moth *Macroglossum stellatarum* (L.) 25.7 - 22.10.95 (2)  
 Elephant Hawk-moth *Deilephila elpenor* (L.) larva 27.9.95 (1) rescued from Blackbird;  
 30.9.95 (1) ? same  
 Pale Prominent *Pterostoma palpina* (Clerck) 29.7.95 (1)  
 Yellow-tail *Euproctis similis* (Fuess.) 1.8 - 3.8.95 (3)  
 Common Footman *Eilema lurideola* (Zinck.) 29.7 - 4.8.95 (28)  
 Garden Tiger *Arctia caja* (L.) 1.8.95 (1)  
 Cream-spot Tiger *Arctia villica* (L.) 8.6 - 13.6.95 (1)  
 Buff Ermine *Spilosoma luteum* (Hufn.) 29.7.95 (2)  
 Ruby Tiger *Phragmatobia fuliginosa* (L.) 6.6 - 28.9.95 (1)  
 White-line Dart *Euxoa tritici* (L.) 29.7 - 1.8.95 (2)  
 Crescent Dart *Agrotis trux* (Hbn.) 29.7 - 4.8.95 (23)  
 Dark Sword-grass *Agrotis ipsilon* (Hufn.) 29.7 - 30.7.95 (3)  
 Flame Shoulder *Ochropleura plecta* (L.) 30.7 - 3.8.95 (1)  
 Large Yellow Underwing *Noctua pronuba* (L.) 29.7 - 4.8.95 (9)  
 Lesser Yellow Underwing *Noctua comes* (Hbn.) 29.7 - 1.8.95 (2)  
 Lesser Broad-bordered Yellow Underwing *Noctua janthina* (D. & S.) 29.7 - 4.8.95 (5)  
 True Lover's Knot *Lycophotia porphyrea* (D. & S.) 29.7 - 1.8.95 (2)  
 Shears *Hada nana* (Hufn.) 30.7 - 4.8.95 (1)  
 Bright-line Brown-eye *Lacanobia oleracea* (L.) 29.7 - 2.8.95 (4)  
 Broom Moth *Ceramica pisi* (L.) 31.7.95 larva (1)  
 Antler *Cerapteryx graminis* (L.) 29.7 - 4.8.95 (8)  
 Clay *Mythimna ferrago* (Fab.) 29.7 - 2.8.95 (1)

Smoky Wainscot *Mythimna impura* (Hbn.) 29.7 - 4.8.95 (3)  
 Common Wainscot *Mythimna pallens* (L.) 29.7.95 (1)  
 Minor Shoulder-knot *Brachyloma viminalis* (Fab.) 29.7 - 30.7.95 (1)  
 Knot Grass *Acronicta rumicis* (L.) 29.7 - 3.8.95 (2)  
 Marbled Green *Cryphia muralis* (Forst.) 1.8.95 (1)  
 Copper Underwing *Amphipyra pyramidea* (L.) 29.7.95 (1)  
 Dark Arches *Apamea monoglypha* (Hufn.) 29.7 - 4.8.95 (17)  
 Rosy Minor *Mesoligia literosa* (Haw.) 29.7 - 4.8.95 (14)  
 Common Rustic *Mesapamea secalis* (L.) 29.7 - 2.8.95 (5)  
 Small Wainscot *Photedes pygmina* (Haw.) 2.8 - 3.8.95 (3)  
 Rustic *Hoplodrina blanda* (D.& S.) 29.7 - 3.8.95 (4)  
 Silver Y *Autographa gamma* (L.) 30.7 - 31.8.95 (33)  
 Spectacle *Abrostola triplasia* (L.) 29.7 - 3.8.95 (2)  
 Snout *Hypena proboscidalis* (L.) 29.7.95 (2)

### FUNGI ON LUNDY 1995

In the dry, summer conditions of 1995, it was extremely difficult to find many fungi, the majority of specimens present being those on dead wood. Samples of all available species were collected by A.J.Parsons and identified by Mr J.G.Keylock, as follows :

*Agrocybe cylindracea* (DC ex Fr.) Maire 4.8.95, on trunk of living sycamore, St Helen's Copse.  
*Bjerkandera adusta* (Willd. ex Fr.) P.Karst 4.8.95, on dead branch of live alder, St Helen's Copse.  
*Bovista nigrescens* (Pers.) ex Pers. 2.8.95, side of track, near Castle.  
*Inonotus radiatus* (Fr.) Karst 3.8.95, on trunk of holm oak, Millcombe; 4.8.95, on alder, St Helen's Copse.  
*Lycogola epidendrum* (L.) Fr. (a myxomycete) 4.8.95, on sycamore and alder, Millcombe and St Helen's Copse.  
*Marasmius rotula* (Fr.) Fr. 4.8.95, on sycamore twigs on ground, St Helen's Copse.  
*Pleurotus ostreatus* (Fr.) Kumm. 31.7.95, on large, dead holly, Millcombe.  
*Polyporus varius* Fr. 29.7.95, on dead gorse branches, St John's Valley.  
*Psathyrella candalleana* (Fr.) Maire 4.8.95, soil at base of alder, St Helen's Copse.  
*Schizopora paradoxa* (Sched. ex Fr.) Domk. 4.8.95, encrusting sycamore branch on ground, Millcombe.  
*Tubaria* agg. 3.8.95, on dead deciduous branches, Millcombe (condition too poor for specific identification).  
*Xylaria polymorpha* (Mérat) Grev. 31.7.95, base of dead sycamore trunk, Millcombe.